Case Report: Cortical Dysplasia

Michael Kean: Michael Ditchfield, M.D.

Children's MRI Centre, Royal Children's Hospital, Parkville, Victoria, Australia

Patient history

10-year-old girl presented after inconclusive scan at an outside institution. The patient has a history of Epilepsy, developmental regression and autistic behavior. The patient was referred to the 3T MAGNETOM Trio and examined using the 32-channel Head Matrix coil utilizing our current high resolution Epilepsy protocol.

Sequence details

The images are from 3T MAGNETOM Trio with software version syngo MR B15 (MPRAGE 0.7 mm isotropic, 0.9 mm isotropic voxels using Water Excitation (WE), DarkFluid 3D SPACE, high resolution 2.5 mm coronal Turbo Spin Echo (TSE) T2 and lipid suppressed 135 ms TE single voxel MR Spectroscopy – MRS)


Image findings

There is a cortically based lesion within the left frontal lobe containing cystic and solid components that do not enhance with contrast. There is a "tail" of increased signal that extends from the lesion to the anterior horn of the lateral ventricle. The MRS confirms appearance of a focal trans mantle cortical dysplasia.

Discussion

The high signal-to-noise ratio (SNR) afforded by the use of the 32-channel Head Matrix coil and 3T permits the technologist to run high-resolution isotropic 3D acquisitions in scan times that are conducive to high patient compliance. The reformatted images (MPRAGE 0.7 mm and 3D SPACE 0.9 mm) combined with the high-resolution T2 provide the radiologist with a level of anatomical detail that provides a diagnosis with a higher degree of sensitivity and specificity. The ability to use these 3D data sets to perform curvilinear reformations improves pre-surgical localization. The lipid suppression characteristics of the MRS


enable lesions close to the outer table of the skull to be evaluated.



1 The reformatted image from 0.7 mm isotropic voxel MPRAGE proves to increase the diagnostic specificity.



2 DarkFluid 3D SPACE shows the left frontal lobe lesion containing solid and cystic components.



3 Lipid suppressed 135 ms single voxel MR spectrum from the lesion of the left frontal lobe supports the diagnosis of a focal trans mantle cortical dysplasia.